arXiv:1203.0895v1 [math.PR] 5 Mar 2012

Smooth-fit principle for a degenerate two-dimensional singular stochastic control problem arising in irreversible investment

Salvatore FEDERICO*
Dipartimento di Economia,
Management e Metodi Quantitativi
Università di Milano
salvatore.federico at unimi.it

Huyên PHAM
Laboratoire de Probabilités et
Modèles Aléatoires, CNRS, UMR 7599
Université Paris Diderot
pham at math.univ-paris-diderot.fr
and CREST-ENSAE

February 24, 2013

Abstract

This paper studies an irreversible investment problem where a social planner aims to control its capacity production in order to fit optimally the random demand of a good. Our model allows for general diffusion dynamics on the demand as well as general cost functional. The resulting optimization problem leads to a degenerate two-dimensional singular stochastic control problem, for which explicit solution is not available in general and the standard verification approach can not be applied a priori. We use a direct viscosity solutions approach for deriving some features of the optimal free boundary function, and for displaying the structure of the solution. In the quadratic cost case, we are able to prove a smooth-fit C^2 property, which gives rise to an explicit identification of the optimal policy and value function.

Keywords: Singular Stochastic Control, Optimal capacity, Irreversible Investment, Viscosity solution, Smooth-fit.

AMS Classification: 93E20, 49J40, 49L25.

---

[*] Part of this research was done when this author was Post-doc at LPMA, Université Paris Diderot, supported by Alma Research.

1 Introduction

We are concerned with a singular stochastic optimal control problem motivated by a model of irreversible investment. More precisely, we imagine to deal with a social planner whose objective is to optimize some functional depending on the current demand of a good (energy, electricity, oil, corn, etc) and its supply in terms of production capacity.

Problems of irreversible investment under uncertainty have been introduced in the economic literature by [23] and then developed by several other authors (see [14, Ch. 11] for references on this subject). From a mathematical point of view, such problems have been formulated as optimal stopping problems or, at a second stage of complexity, as singular stochastic optimal control problems, and have given a considerable impulse to the development of the corresponding mathematical theory. As references for the theory of singular stochastic control in context different from irreversible investment, we may mention the works [12, 19, 20, 21] and [15, Ch. VIII]. The mathematical literature of singular stochastic control applied to the subject of irreversible investment under uncertainty includes the works [3, 4, 5, 6, 10, 11, 13, 25, 28, 31]. The economic issue of costly reversibility has then been introduced and studied, among others, in [1, 4, 16, 17, 22, 24]. In the papers mentioned above, the ones (substantially) considering two state variables (an uncontrolled one - possibly multidimensional - containing the noise, and a controlled one, representing the capacity) are [3, 4, 17, 22, 24, 28, 31][1]. More precisely, the ones dealing with a dynamic programming approach directly on the singular control problem and with the study of the associated Hamilton-Jacobi-Bellman equation (which in this case is a variational inequality) are [22, 24, 31]. In particular, the papers [24, 31] consider an expected performance on infinite horizon with discounting over time, as in our case. However, the approach of [24, 31] is of *verification* type. In a singular stochastic control framework, this means that one has to guess some smooth fit properties of the value function at the optimal free boundary in order to look for a solution of the Hamilton-Jacobi-Bellman equation. Then one needs to prove *a posteriori* that the solution found is indeed the value function and, as a byproduct, one gets also the optimal feedback control. When this approach is applicable, it turns out to be very convenient, as it is theoretically fast (even if it may involve a nontrivial technical complexity) and allows a first understanding of the problem. Moreover, the presence of an explicit solution is an important tool to analyze the qualitative properties of optimal control and trajectory. On the other hand, one has to recognize that it presents two drawdowns. First, it is based on a guess, and so it cannot bring to a deep understanding of the structural background of the problem. Second, it works only when explicit solutions are available, therefore it leaves the problem completely unsolved most of the cases.

In the present paper, we perform a direct study of the problem (without passing through verification type arguments) by means of a viscosity approach to the Hamilton-Jacobi-Bellman

---

[1] The paper [3] reduces first the singular control problem to an optimal stopping problem and then applies dynamic programming to the latter one. The paper [17] reduces first the singular control problem to an optimal switching problem and then applies dynamic programming to the latter one. Finally, we should mention the paper [30], dealing with a singular control problem with two state variables in a different context (consumption-investment under transaction costs). In this case the problem is approached by dynamic programming and by means of viscosity solutions to the associated Hamilton-Jacobi-Bellman equation. However, the regularity of the value function is proved by reducing the problem to dimension one, which is possible in that case due to the specific structure of the problem.

(HJB) equation. To our knowledge, this is the first time that such an approach is used in the case of two state variables, in particular when the controlled state variable, here the capacity process, has no diffusion term, and so is degenerate[2]. Our approach allows us to keep much more generality with regard to the uncontrolled state variable (which is indeed a very general diffusion in the present paper) and to state the smooth-fit conditions of [24] as *necessary* conditions of optimality, i.e. prove that the value function *must* satisfy these conditions[3]. More precisely, we show that the value function is C^1 along the component of the controlled variable (Proposition 3.1; this easily follows from our assumptions by convexity arguments, just working on the definition of value function). Then, we prove that it has continuous mixed second derivative along the optimal boundary function (Proposition 5.1 and Remark 5.1; this is a deeper result, which invokes the viscosity property of the value function and requires the additional assumption (5.4)). At the end, this machinery allows us to uniquely individuate the value function by Theorems 4.1 and 5.1, and solve the problem by Theorem 4.2.

The rest of the paper is organized as follows. In Section 2, we formulate the two-dimensional singular stochastic control problem and state the main assumptions. We study in Section 3 some first properties of the value function and of the optimal boundary, which is a function of the demand. In Section 4, by relying on the viscosity property of the value function to its dynamic programming variational inequality, we give a first main result providing the structure of the value function, and state a second main result yielding the optimal control in terms of the optimal boundary. Section 5 focus on the case of quadratic cost function, which allows us to prove a second order smooth fit principle. This leads to the missing information to explicitly individuate the value function and the optimal boundary (the third main result), and makes the results of Section 4 applicable. Finally, we close the paper by explicit illustrations of the theory to the basic example of geometric Brownian motion for the uncontrolled demand diffusion. More examples and applications are developed in the companion paper [2] where we also take into account delay in the expansion of the capacity production.

2 The singular stochastic control problem

Let us fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ equipped with a filtration $\mathbb{F} = (\mathcal{F}_t)_{t\geq 0}$ satisfying the usual conditions, and supporting a standard one-dimensional Brownian motion $(W_t)_{t\geq 0}$. On this space, we consider an uncontrolled state process $D = (D_t)_{t\geq 0}$ (representing the demand of a good), governed by a regular diffusion dynamics:

---

[2] There are of course several papers, which consider singular stochastic control problems with multidimensional state variables, and characterize the value function in terms of viscosity solutions to the associated HJB equations. However, rather few go beyond the viscosity characterization, and investigate smooth-fit properties in order to derive the structural form of the value function. In this spirit, we may mention the paper [16] in the case of just one dimensional controlled variable. See also [18] for impulse control of multi-dimensional diffusion processes with non degenerate diffusion term. On the other hand, we may quote the paper [29], which studies regularity of a two-dimensional singular control problem with nondegenerate diffusion.

[3] Another major advantage of such approach is that it allows generalizations. With this regard, we notice that here we minimize a cost functional and that we are in an irreversible investment environment. However, the arguments used here can be extended to the case of profit/cost functional and reversibility of investment, as in [24, 31].

$$\mathrm{d}D_t \;=\; \mu(D_t)\mathrm{d}t + \sigma(D_t)\mathrm{d}W_t, \quad D_0 = d. \tag{2.1}$$

Throughout the paper we assume the following.

Assumption 2.1. *(i) The coefficients μ, σ are continuous, have at most linear growth, i.e. there exist constants C_μ, C_σ such that*

$$|\mu(d)| \;\leq\; C_\mu(1 + |d|), \quad |\sigma(d)| \;\leq\; C_\sigma(1 + |d|), \quad \forall d \in \mathbb{R}, \tag{2.2}$$

and are such that there exists a unique solution to (2.1) *on* $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0}, \mathbb{P})$.
(ii) The solution takes values in some open interval

$$\mathcal{O} \;=\; (d_{min}, d_{max}), \quad -\infty \leq d_{min} < d_{max} \leq +\infty.$$

with not-exit boundaries d_{min}, d_{max} (in the sense of Feller's classification, see [7, Ch. II]). Moreover, we also assume that the boundary d_{min} is not-entrance (and consequently natural, see again [7, Ch. II]).
(iii) The diffusion is nondegenerate, i.e. $\sigma^2(d) > 0$ for all $d \in \mathcal{O}$.

Next, we consider a control process $I = (I_t)_{t\geq 0}$, which is a càdlàg nondecreasing $\mathbb{F}$-adapted process. I_t represents the cumulative irreversible investment done from time 0 up to time $t \geq 0$. We denote by $\mathcal{I}$ the set of such control processes I s.t. $I_{0^-} = 0$. The production capacity process $(C_t)_{t\geq 0}$, controlled by $I \in \mathcal{I}$, is given by

$$C_t \;=\; c + I_t, \quad C_{0^-} = c \in \mathbb{R}. \tag{2.3}$$

The objective is to minimize over $I \in \mathcal{I}$ the functional

$$\mathbb{E}\Big[\int_0^\infty e^{-\rho t}\Big(g(C_t, D_t)\mathrm{d}t + q_0\mathrm{d}I_t\Big)\Big], \tag{2.4}$$

where $g : \mathbb{R} \times \mathcal{O} \to [0, \infty)$ is a running cost function, $q_0 > 0$ is the cost per unit of investment, and ρ is a positive discount factor. Notice that with respect to the usual irreversible investment literature, which is mainly based on profit/cost performance criterions, we focus here on the minimization of a cost criterion in the spirit of a social planning problem, whose objective is to fit the capacity production to the demand at cheapest cost, see [2] for a detailed description and explanation. Actually, we shall make the following assumptions on the cost function g.

Assumption 2.2. *(i) g is continuous on $\mathbb{R} \times \mathcal{O}$, $g(\cdot, d)$ is strictly convex for all $d \in \mathcal{O}$, and α-semiconcave with $\alpha \in (0, 1]$ uniformly with respect to $d \in \mathcal{O}$, meaning that there exists $M_0 > 0$ such that for every $\lambda \in [0, 1]$, $c, c' \in \mathbb{R}$ and $d \in \mathcal{O}$,*

$$\lambda g(c, d) + (1 - \lambda)g(c', d) - g(\lambda c + (1 - \lambda)c', d) \;\leq\; M_0\lambda(1 - \lambda)|c - c'|^{1+\alpha}. \tag{2.5}$$

(ii) The partial derivative function $g_c(c, \cdot)$ is nonincreasing in $\mathcal{O}$ for every $c \in \mathbb{R}$.
(iii) g satisfies a polynomial growth condition: there exist $C_0 > 0$ $\nu \geq 0$, such that

$$g(c, d) \;\leq\; C_0(1 + |c|^\nu + |d|^\nu), \quad \forall\, c \in \mathbb{R},\; d \in \mathcal{O}.$$

Remark 2.1. 1. The α-semiconcavity property (2.5) is verified if and only if $g(\cdot, d) \in C^{1+\alpha}(\mathbb{R}; \mathbb{R}^+)$ uniformly in $d \in \mathcal{O}$, in the sense that $g(\cdot, d) \in C^1(\mathbb{R}; \mathbb{R})$ for every $d \in \mathcal{O}$ and and $g_c(\cdot, d)$ is α-Hölder continuous uniformly in $d \in \mathcal{O}$. This is consequence of Propositions 2.1.2 and 2.1.3 in [9].
2. We observe that Assumption 2.2-(ii) has an economic interpretation. It means that the marginal cost with respect to capacity for a fixed level of capacity is nonincreasing in the demand: since this marginal cost measures the instantaneous convenience to invest a unit of capacity, this assumption says that if the demand is going to increase, it becomes more convenient to invest.
3. Any function g of the spread $|c - d|$ between capacity and demand, in the form

$$g(c, d) \quad = \quad C_0 |c - d|^{1+\alpha}, \quad C_0 \geq 0, \ \alpha \in (0, 1], \tag{2.6}$$

satisfies Assumption 2.2. □

Remark 2.2. Following the idea of [5, Sec. 6], our model admits a suitable generalization to the case of capacity dynamics in the form:

$$\mathrm{d}C_t \quad = \quad C_t (b \, \mathrm{d}t + \gamma \, \mathrm{d}W_t^0) + \mathrm{d}I_t, \ \ C_{0^-} \ = \ c,$$

where W^0 is another Browinan motion independent of W. Indeed letting C^0 be the solution to

$$\mathrm{d}C_t^0 \quad = \quad C_t^0 (b \, \mathrm{d}t + \gamma \, \mathrm{d}W_t^0), \ \ C_0^0 \ = \ 1,$$

the process C can be rewritten as

$$C_t \ = \ C_t^0 [c + \bar{I}_t], \quad \text{with} \quad \bar{I}_t \ = \ \int_0^t \frac{1}{C_s^0} \, \mathrm{d}I_s,$$

and the problem becomes

$$\inf_{\bar{I} \in \mathcal{I}} \mathbb{E} \Big[\int_0^\infty e^{-\rho t} \Big(g(C_t^0 \, [c + \bar{I}_t], D_t) + q_0 C_t^0 \mathrm{d}\bar{I}_t \Big) \Big].$$

This problem involves an additional uncontrolled state variable (the variable C^0), but keeps the basic structures, so it seems approachable by the same techniques developed in the next sections. □

3 Some properties of the optimal boundary

We shall study the optimization problem (2.4) by dynamic programming methods, and so consider this singular stochastic control problem when varying initial data. Therefore, from now on, we stress the dependence of C on c, I and the dependence of D on d by denoting them respectively as $C^{c,I}$, D^d. Although the economic problem is meaningful for positive c, we also allow for mathematical convenience negative values of c. The state space is then equal to

$$\mathcal{S} \quad = \quad \mathbb{R} \times \mathcal{O}.$$

Given $(c, d) \in \mathcal{S}$, the functional to be minimized over $I \in \mathcal{I}$ is

$$G(c, d; I) \;=\; \mathbb{E}\Big[\int_0^\infty e^{-\rho t}\Big(g(C_t^{c,I}, D_t^d)\mathrm{d}t + q_0\mathrm{d}I_t\Big)\Big],$$

and the associated value function is

$$v(c, d) \;=\; \inf_{I \in \mathcal{I}} G(c, d; I), \quad (c, d) \in \mathcal{S}. \tag{3.1}$$

Notice that $v \geq 0$ as $g \geq 0$. On the other hand, taking the null control $I \equiv 0$, we have an upper bound for the value function:

$$v(c, d) \;\leq\; \hat{V}(c, d) \;:=\; \mathbb{E}\Big[\int_0^\infty e^{-\rho t} g(c, D_t^d)\mathrm{d}t\Big]. \tag{3.2}$$

Under (2.2), and by standard estimates, we know that that there exist positive constants $K_0 = K_{0,\mu,\sigma,\nu}$, $K_1 = K_{1,\mu,\sigma,\nu}$ such that

$$\mathbb{E}\Big[\big|D_t^d\big|^\nu\Big] \;\leq\; K_0(1 + |d|^\nu)e^{K_1 t}, \quad \forall t \geq 0. \tag{3.3}$$

In the sequel, we make the standing assumption that the discount factor ρ satisfies

$$\rho \;>\; K_1, \tag{3.4}$$

where K_1 is the constant appearing in (3.3). It is then easy to check with (3.3) that $\hat{V}$, and hence also v inherits from g the polynomial growth condition in Assumption 2.2-(iii): there exists $C_0 = C_{0,\mu,\sigma,\nu}$ s.t.

$$0 \;\leq\; v(c, d) \;\leq\; \hat{V}(c, d) \;\leq\; C_0(1 + |c|^\nu + |d|^\nu), \quad \forall (c, d) \in \mathcal{S}, \tag{3.5}$$

which implies in particular that the value function v is finite and locally bounded.

We next show directly some preliminary regularity properties on $\hat{V}$ and on the value function.

Lemma 3.1. *The function $\hat{V}$ belongs to the class $C^{1+\alpha,2}(\mathcal{S}; \mathbb{R})$, meaning that it is once differentiable with respect to c with α-Hölder continuous derivative, and twice differentiable with respect to d. Moreover, $\hat{V}_c$ is continuous on $\mathcal{S}$.*

Proof. First, we notice by standard arguments that $\hat{V}(\cdot, d)$ is convex. Now, let $\lambda \in [0, 1]$, $c, c' \in \mathbb{R}$, $d \in \mathcal{O}$, and set $c_\lambda = \lambda c + (1 - \lambda)c'$. Recalling Assumption 2.2-(i), we have

$$\begin{aligned}
& \lambda\hat{V}(c, d) + (1 - \lambda)\hat{V}(c', d) - \hat{V}(c_\lambda, d) \\
= \;& \mathbb{E}\Big[\int_0^\infty e^{-\rho t}\Big(\lambda g(c, D_t^d) + (1 - \lambda)g(c', D_t^d) - g(c_\lambda, D_t^d)\Big)\,\mathrm{d}t\Big] \\
\leq \;& M_0\lambda(1 - \lambda)\int_0^\infty e^{-\rho t}|c - c'|^{1+\alpha}\mathrm{d}t \;=\; \frac{M_0}{\rho}\lambda(1 - \lambda)|c - c'|^{1+\alpha},
\end{aligned}$$

which shows that $\hat{V}$ is α-semiconcave in c (uniformly w.r.t. d). Together with the convexity of $\hat{V}(., d)$, and by Proposition 2.1.3 in [9], this shows that $\hat{V}$ is once differentiable with respect to c with α-Hölder continuous derivative. Twice differentiability with respect to d is a consequence of the fact that the diffusion is nondegenerate, as by standard theory $\hat{V}(c,\cdot)$ solves a uniformly elliptic equation on $\mathcal{O}$.

Moreover, by dominated convergence, it is possible to differentiate under the integral sign in (3.2) and get the representation

$$\hat{V}_c(c,d) \;=\; \mathbb{E}\Big[\int_0^\infty e^{-\rho t} g_c(c, D_t^d)\mathrm{d}t\Big].$$

Then, by similar arguments, one obtain the twice differentiability of $\hat{V}_c$ with respect to d. Finally, continuity of V_c in the couple (c,d) is consequence of its local Lipschitz continuity with respect to d and its α-Hölder continuity with respect to c. □

Proposition 3.1. *The value function v is convex with respect to c and α-semiconcave with respect to c uniformly in $d \in \mathcal{O}$. Hence, $v(\cdot,d) \in C^{1+\alpha}(\mathbb{R};\mathbb{R})$ uniformly in $d \in \mathcal{O}$.*

Proof. Convexity of v follows from the convexity of g with respect to c and linearity of the state equation for $C^{c,I}$. Let us now prove the semiconcavity property. Let $\varepsilon > 0$, $\lambda \in [0,1]$, $c, c' \in \mathbb{R}$, $d \in \mathcal{O}$, set $c_\lambda = \lambda c + (1-\lambda)c'$, and take an ε-optimal control $I^\varepsilon \in \mathcal{I}$ for (c_λ, d). Due to Assumption 2.2-(i), we have

$$\begin{aligned}\lambda v(c,d) + (1-\lambda)v(c',d) - v(c_\lambda,d) &\leq \lambda G(c,d;I^\varepsilon) + (1-\lambda)G(c',d;I^\varepsilon) - G(c_\lambda,d;I^\varepsilon) + \varepsilon \\ &\leq M_0\lambda(1-\lambda)\int_0^\infty e^{-\rho t}|c-c'|^{1+\alpha}\mathrm{d}t + \varepsilon \\ &= \frac{M_0}{\rho}\lambda(1-\lambda)|c-c'|^{1+\alpha} + \varepsilon.\end{aligned}$$

By arbitrariness of ε, this shows the required semiconcavity property. The last claim follows by convexity and α-semiconcavity of $v(\cdot,d)$, by using again Proposition 2.1.3 in [9]. □

Proposition 3.2. *The following properties hold true:*

(i) $v_c(\cdot,d) \geq -q_0$ *for each* $d \in \mathcal{O}$.

(ii) $v_c(c,\cdot)$ *is nonincreasing for each* $c \in \mathbb{R}$.

Proof. (i) Let $(c,d) \in \mathcal{S}$ and, for each $I \in \mathcal{I}$, $\varepsilon > 0$, set $I^\varepsilon := I + \varepsilon$. Then we have

$$G(c,d;I) - v(c-\varepsilon,d) \;\geq\; G(c,d;I) - G(c-\varepsilon,d;I^\varepsilon) \;=\; -q_0\varepsilon.$$

Taking the infimum over $I \in \mathcal{I}$, dividing by ε and letting $\varepsilon \to 0$, we get the required inequality for v_c.

(ii) Fix $(c,d') \in \mathcal{C}$ and set $\alpha = v_c(c,d')$. Then for every $\delta > 0$ there exist $\varepsilon > 0$ such that

$$\frac{v(c,d') - v(c-\varepsilon,d')}{\varepsilon} \;\geq\; \alpha - \delta.$$

By taking $I^{\varepsilon\delta} \in \mathcal{I}$ an $\varepsilon\delta$-optimal control for $(c-\varepsilon,d')$, we then get

$$G(c,d';I) - G(c-\varepsilon,d';I^{\varepsilon\delta}) \;\geq\; \varepsilon(\alpha - 2\delta), \quad \forall I \in \mathcal{I}. \tag{3.6}$$

Now, under Assumption 2.2-(ii), we have

$$g(c,d) - g(c',d) \quad \geq \quad g(c,d') - g(c',d'), \quad \forall c,c' \in \mathbb{R}, \ \forall d \leq d',$$

which combined with (3.6) yields

$$G(c,d;I) - G(c-\varepsilon,d;I^{\varepsilon\delta}) \quad \geq \quad \varepsilon(\alpha - 2\delta), \quad \forall I \in \mathcal{I}, \ \forall d \leq d',$$

and so:

$$G(c,d;I) - v(c-\varepsilon,d) \quad \geq \quad \varepsilon(\alpha - 2\delta), \quad \forall I \in \mathcal{I}, \ \forall d \leq d'. \tag{3.7}$$

Taking the infimum on $I \in \mathcal{I}$ in (3.7), we then get

$$v(c,d) - v(c-\varepsilon,d) \quad \geq \quad \varepsilon(\alpha - 2\delta), \quad \forall d \leq d',$$

which shows by convexity of $v(\cdot,d)$ that

$$v_c(c,d) \quad \geq \quad (\alpha - 2\delta), \quad \forall d \leq d'.$$

By arbitrariness of δ, this means $v_c(c,d) \geq v_c(c,d')$, for all $d \leq d'$, which ends the proof. $\square$

In view of Proposition 3.2 (i), we introduce the so-called *continuation region*

$$\mathcal{C} \quad := \quad \{(c,d) \in \mathcal{S} \mid v_c(c,d) > -q_0\},$$

and its complement set, the *action region*

$$\mathcal{A} \quad := \quad \{(c,d) \in \mathcal{S} \mid v_c(c,d) = -q_0\}. \tag{3.8}$$

Let us then consider the function $\hat{c} : \mathcal{O} \to \bar{\mathbb{R}}$ defined by

$$\hat{c}(d) \quad := \quad \inf\{c \in \mathbb{R} \mid v_c(c,d) > -q_0\},$$

with the convention $\inf \emptyset = \infty$, $\inf \mathbb{R} = -\infty$. Actually, since $v(\cdot,d)$ is convex and nonnegative, we notice that $\hat{c}(d) < \infty$ for all $d \in \mathcal{O}$. Then, by convexity of v with respect to c, the action and continuation regions are expressed in terms of the free boundary function $\hat{c}$ as:

$$\mathcal{C} \ = \ \{(c,d) \in \mathcal{S} \mid c > \hat{c}(d)\}, \qquad \mathcal{A} \ = \ \{(c,d) \in \mathcal{S} \mid c \leq \hat{c}(d)\}. \tag{3.9}$$

We denote by $\mathring{\mathcal{C}}$ the topological interior of $\mathcal{C}$ (at this stage we do not know yet that $\mathcal{C}$ is open) and by $\bar{\mathcal{C}}$ its topological closure in $\mathcal{S}$. It is clear that

$$\bar{\mathcal{C}} \quad \supset \quad \{(c,d) \in \mathcal{S} \mid c \geq \hat{c}(d)\}. \tag{3.10}$$

We also introduce the functions $\hat{c}_g$ and $\hat{c}_{\hat{V}}$ from $\mathcal{O}$ into $[-\infty,\infty)$ defined by:

$$\hat{c}_g(d) \ = \ \inf\{c \in \mathbb{R} \mid g_c(c,d) > -\rho q_0\}, \qquad \hat{c}_{\hat{V}}(d) \ = \ \inf\{c \in \mathbb{R} \mid \hat{V}_c(c,d) > -q_0\}.$$

One easily checks that they are nondecreasing (by Assumption 2.2-(ii)) and right-continuous (by continuity of $g_c, \hat{V}_c$).

Proposition 3.3. *The function $\hat c$ is nondecreasing and $\hat c_{\hat V} \le \hat c \le \hat c_g$.*

Proof. *Monotonicity.* The nondecreasing property of $\hat c$ is a consequence of the nonincreasing monotonicity of $d \mapsto v_c(c,d)$ (Proposition 3.2-(ii)).

Estimate from above. Let $d \in \mathcal{O}$ and $c > \hat c_g(d)$. Then there exists $\varepsilon > 0$ such that $c - \varepsilon > \hat c_g(d)$. By definition and right-continuity of $\hat c_g$, and by Assumption 2.2 (ii), there exist $\bar d > d$ and $\delta > 0$ such that

$$g_c(\zeta,\eta) \;\; \ge \;\; -\rho q_0 + \delta, \quad \forall \zeta > c-\varepsilon,\ \forall \eta < \bar d. \tag{3.11}$$

Let us consider the stopping time

$$\tau \;\; = \;\; \inf\{t \ge 0 \mid D^d_t \ge \bar d\} \;\; > \;\; 0 \;\; a.s.$$

By (3.11) and the mean value theorem, we then have

$$g(C^{c,I}_t, D^d_t) - g(c-\varepsilon, D^d_t) \;\; \ge \;\; (-\rho q_0 + \delta)(C^{c,I}_t - c + \varepsilon), \quad \forall t \in [0,\tau). \tag{3.12}$$

Given an arbitrary $I \in \mathcal{I}$, define $I^\varepsilon_t = (I_t + \varepsilon)1_{t \ge \tau}$, and notice that $C^{c,I}_t = C^{c-\varepsilon, I^\varepsilon}_t$, for all $t \ge \tau$. We then have

$$\begin{aligned}
& G(c,d;I) - G(c-\varepsilon,d;I^\varepsilon) \\
= \;\; & \mathbb{E}\Big[\int_0^\tau e^{-\rho t}\Big(g(C^{c,I}_t, D^d_t)\mathrm{d}t + q_0 \mathrm{d}I_t\Big)\Big] - \mathbb{E}\Big[\int_0^\tau e^{-\rho t} g(c-\varepsilon, D^d_t)\mathrm{d}t + e^{-\rho\tau} q_0 (I_\tau + \varepsilon)\Big] \\
= \;\; & \mathbb{E}\Big[\int_0^\tau e^{-\rho t}\Big(g(C^{c,I}_t, D^d_t) - g(c-\varepsilon, D^d_t)\Big)\mathrm{d}t\Big] + \mathbb{E}\Big[\int_0^\tau e^{-\rho t} q_0 \mathrm{d}I_t - e^{-\rho\tau} q_0 (I_\tau + \varepsilon)\Big] \\
= \;\; & \mathbb{E}\Big[\int_0^\tau e^{-\rho t}\Big(g(C^{c,I}_t, D^d_t) - g(c-\varepsilon, D^d_t)\Big)\mathrm{d}t\Big] + \mathbb{E}\Big[\rho q_0 \int_0^\tau e^{-\rho t} I_t \mathrm{d}t - e^{-\rho\tau} q_0 \varepsilon\Big],
\end{aligned} \tag{3.13}$$

where we used integration by parts in the last equality. By plugging (3.12) into (3.13), and since $I_t = C^{c,I}_t - c$, we get

$$G(c,d;I) - G(c-\varepsilon,d;I^\varepsilon) \;\; \ge \;\; -\varepsilon q_0 + \varepsilon\delta\, \mathbb{E}\Big[\int_0^\tau e^{-\rho t}\mathrm{d}t\Big],$$

and so

$$G(c,d;I) - v(c-\varepsilon,d) \;\; \ge \;\; (-q_0 + \delta')\varepsilon, \tag{3.14}$$

where we set $\delta' = \delta \mathbb{E}\big[\int_0^\tau e^{-\rho t}\mathrm{d}t\big] > 0$. Taking the infimum over $I \in \mathcal{I}$ in (3.14), and dividing by ε, we get

$$\frac{v(c,d) - v(c-\varepsilon,d)}{\varepsilon} \;\; \ge \;\; -q_0 + \delta'$$

which combined with the convexity of $v(\cdot,d)$, shows that $v_c(c,d) > -q_0$. This proves that $c > \hat c(d)$, and so the required upper bound for $\hat c(d)$.

Estimate from below. Recall by convexity of $g(\cdot,d)$ that

$$g(c'+r,d) - g(c',d) \;\; \ge \;\; g(c+r,d) - g(c,d), \quad \forall c' \ge c,\ \forall r \ge 0,\ \forall d \in \mathcal{O}. \tag{3.15}$$

Let $d \in \mathcal{O}$ and $c > \hat{c}(d)$. Then by definition of $\hat{c}(d)$ and convexity of $v(\cdot, d)$, there exist $\varepsilon, \delta > 0$ such that

$$v(c,d) - v(c-\varepsilon, d) \quad \geq \quad \varepsilon(-q_0 + 2\delta).$$

Taking a control $I^{\varepsilon\delta} \in \mathcal{I}$, which is $\varepsilon\delta$-optimal for $v(c-\varepsilon, d)$, we then obtain:

$$G(c,d;I^{\varepsilon\delta}) - G(c-\varepsilon, d; I^{\varepsilon\delta}) \quad \geq \quad \varepsilon(-q_0 + \delta). \tag{3.16}$$

Now observe that $C_t^{c,I^{\varepsilon\delta}} = c + I_t^{\varepsilon\delta}$, $C_t^{c-\varepsilon,I^{\varepsilon\delta}} = c - \varepsilon + I_t^{\varepsilon\delta}$. Then, from (3.15) and (3.16), we get

$$\hat{V}(c,d) - \hat{V}(c-\varepsilon, d) \quad \geq \quad \varepsilon(-q_0 + \delta).$$

Since $\hat{V}(\cdot, d)$ is convex, this shows that $\hat{V}_c(c,d) > -q_0$, which yields $c > \hat{c}_{\hat{V}}(d)$, and the claim is proved. $\square$

Let us define

$$\underline{c} \; := \; \inf_{d \in \mathcal{O}} \hat{c}(d), \qquad \bar{c} \; := \; \sup_{d \in \mathcal{O}} \hat{c}(d),$$

and the pseudo-inverse function $\hat{d}$ of $\hat{c}$ defined on $\mathbb{R}$ by

$$\hat{d}(c) \quad := \quad \inf\{d \in \mathcal{O} \mid v_c(c,d) = -q_0\}, \tag{3.17}$$

with the convention $\inf \emptyset = d_{max}$. We also introduce the c-section sets of the continuation region

$$S_c \quad := \quad \{(c,d) \mid d \in (d_{min}, \hat{d}(c))\}, \quad c \in \mathbb{R}.$$

We have the following result on the form of the continuation region.

Proposition 3.4. *(1) The function $\hat{d}$ is nondecreasing and $\lim_{c\to\infty} \hat{d}(c) = d_{max}$.*

(2) S_c is increasing in c, in the sense that $S_c \subset S_{c'}$ if $c \leq c'$.

(3) $S_c \neq \emptyset$ if and only if $c > \underline{c}$.

(4) If $c > \bar{c}$ then $S_c = (d_{min}, d_{max})$, and if $S_c = (d_{min}, d_{max})$ then $c \geq \bar{c}$.

(5) We have the representation

$$\overset{\circ}{\mathcal{C}} \quad = \quad \bigcup_{c \in \mathbb{R}} S_c. \tag{3.18}$$

Proof. (1) The fact that $\hat{d}$ is nondecreasing follows from monotonicity of v_c with respect to d. By monotonicity of $\hat{d}$, the limit $\lim_{c\to+\infty} \hat{d}(c)$ exists. Suppose by contradiction that $\lim_{c\to+\infty} \hat{d}(c) = \bar{d} < d_{max}$. Then, by monotonicity of v_c with respect to d, we have $v_c(c,d) = -q_0$ for each $(c,d) \in \mathbb{R} \times (\bar{d}, d_{max})$. But this is not possible since v is nonnegative, so we have the claim. The assertion (2) is a direct consequence of the monotonicity of $\hat{d}$, while assertions (3) and (4) follow from the definition of $\underline{c}$ and $\bar{c}$. Let us prove the last assertion (5). If $(c,d) \in \overset{\circ}{\mathcal{C}}$, then $\hat{v}_c > -q_0$ in some suitable neighborhood of (c,d). Therefore, by definition of $\hat{d}(c)$ and monotonicity of v_c with respect to d, this implies that $(c, d) \in S_c \neq \emptyset$ for some $c \in \mathbb{R}$. Hence, the inclusion $\overset{\circ}{\mathcal{C}} \subset \bigcup_{c \in \mathbb{R}} S_c$ is proved. Conversely, let (c, d) be such that $(c, d) \in S_c$ for some $c \in \mathbb{R}$. By assertion (3), we notice that $c > \underline{c}$. By definition of $\hat{d}$, we get: $v_c(c, \cdot) > -q_0$ in some neighborhood of d. Together with the continuity of v_c with respect to c, and the monotonicity of v_c with respect to d, this shows actually that the inequality $v_c > -q_0$ holds in some neighborhood of (c, d). This proves the inclusion $\overset{\circ}{\mathcal{C}} \supset \bigcup_{c \in \mathbb{R}} S_c$, and finally the required relation (3.18). □

4 Structure of the solution

The dynamic programming equation for the singular stochastic control (3.1) takes the form of a variational inequality:

$$\max\big\{[\mathcal{L}v(c, \cdot)](d) - g(c, d), -v_c(c, d) - q_0\big\} \quad = \quad 0, \quad (c, d) \in \mathcal{S}, \tag{4.1}$$

where the second-order differential operator $\mathcal{L}$ is defined by

$$[\mathcal{L}\varphi](d) \quad = \quad \rho\varphi(d) - \mu(d)\varphi'(d) - \frac{1}{2}\sigma(d)^2\varphi''(d).$$

In the following, given a locally bounded function $\phi : \mathbb{R} \to \mathbb{R}$, we denote respectively by φ^*, and φ_* the upper semicontinuous and the lower semicontinuous envelope of ϕ. Since we do not know a priori if there exists a smooth solution to (4.1), we first rely in general on the notion of viscosity solutions, whose definition in our context is given by:

Definition 4.1. *(i) We say that v is a viscosity subsolution to* (4.1) *if for any* $(c, d) \in \mathcal{S}$,

$$\max\big\{[\mathcal{L}\varphi(c, \cdot)](d) - g(c, d), -\varphi_c(c, d) - q_0\big\} \quad \leq \quad 0,$$

whenever $\varphi(c, .) \in C^2(\mathbb{R}; \mathbb{R})$ *and* $v(c, \cdot)^* - \varphi(c, \cdot)$ *has a local maximum at* d.
(ii) We say that v is a viscosity supersolution to (4.1) *if for any* $(c, d) \in \mathcal{S}$,

$$\max\big\{[\mathcal{L}\varphi(c, \cdot)](d) - g(c, d), -\varphi_c(c, d) - q_0\big\} \quad \geq \quad 0,$$

whenever $\varphi(c, .) \in C^2(\mathbb{R}; \mathbb{R})$ *and* $v(c, \cdot)_* - \varphi(c, \cdot)$ *has a local minimum at* d.
(iii) We say that v is a viscosity solution to (4.1) *if is is both a viscosity sub- and supersolution.*

The viscosity property of the value function follows usually from the dynamic programming principle (DPP). The statement of DPP calls upon delicate measurable selection arguments. Once we know a priori that the value function is continuous, measurable selection holds true and follows by classical arguments, see e.g. [15]. However, since we are not assuming Lipschitz continuity of the coefficients in (2.1), is it not clear how to get the continuity of the value function from its very definition. Instead, we can use the concept of weak dynamic programming introduced in [8], which holds for our problem, and then derive by standard arguments of stochastic control (see e.g. [26, Ch. 4]) the viscosity property of the value function.

Proposition 4.1. *The value function v is a viscosity solution to* (4.1) *on* $\mathcal{S}$.

Remark 4.1. One could prove a comparison principle to the variational inequality (4.1) for viscosity sub-and super solution satisfying the growth condition (3.5), hence providing a uniqueness viscosity characterization of the value function v. However, we can avoid such comparison principle issue in our approach relying mainly on the viscosity property in order to derive a smooth-fit property, which then allows us to identify uniquely the value function. □

Remark 4.2. Since the value function v is continuously differentiable w.r.t. the argument c (Proposition 3.1), the viscosity property of v to (4.1) implies in particular the following viscosity property on the continuation region (where $v_c > -q_0$):

$$[\mathcal{L}v(c,\cdot)](d) - g(c,d) \;=\; 0, \qquad (c,d) \in \mathcal{C}. \tag{4.2}$$

□

We now investigate the structure of the value function v in the continuation region $\mathcal{C}$ and in the action region $\mathcal{A}$, and state some topological properties of these sets. Let us denote respectively by ψ and φ the strictly positive, increasing and strictly positive, decreasing fundamental solutions to the linear ordinary differential equation

$$\mathcal{L}\phi(d) \;:=\; \rho\phi(d) - \mu(d)\phi'(d) - \frac{1}{2}\sigma(d)^2\phi''(d) \;=\; 0,$$

which satisfy

$$\psi(d_{min}^{+}) \;=\; 0, \qquad \varphi(d_{min}^{+}) \;=\; \infty, \tag{4.3}$$

since d_{min} is a natural boundary. The existence and properties of such functions can be found in several references including Chapter 2 in [7].

Lemma 4.1. *Let $c > \underline{c}$, and $\hat{d}$ defined by* (3.17)*. Then*

$$v(c,\cdot) \;\in\; C^0((d_{min},\hat{d}(c)];\mathbb{R}) \;\cap\; C^2((d_{min},\hat{d}(c));\mathbb{R}).$$

Moreover there exist constants $A(c), B(c) \in \mathbb{R}$ such that

$$v(c,d) \;=\; A(c)\psi(d) + B(c)\varphi(d) + \hat{V}(c,d), \quad d \in (d_{min},\hat{d}(c)]. \tag{4.4}$$

Proof. Fix $c > \underline{c}$ and $[a,b] \subset (d_{min},\hat{d}(c)]$, and consider the Dirichlet problem

$$\begin{cases} \rho u(d) - \mu(d)u'(d) - \frac{1}{2}\sigma(d)^2 u''(d) = g(c,d), & d \in (a,b), \\ u(a) = v(c,a), \quad u(b) = v(c,b). \end{cases} \tag{4.5}$$

This problem admits a unique viscosity solution, which must coincide with $v(c,\cdot)$ in $[a,b]$ from (4.2). On the other hand, by uniform ellipticity of σ, problem (4.5) admits a solution of class $C^0([a,b];\mathbb{R}) \cap C^2((a,b);\mathbb{R})$. Hence, we deduce that $v(c,\cdot) \in C^0((d_{min},\hat{d}(c)];\mathbb{R}) \cap C^2((d_{min},\hat{d}(c));\mathbb{R})$, and satisfies in a classical sense:

$$[\mathcal{L}v(c,\cdot)](d) - g(c,d) \;=\; 0, \qquad d \in (d_{min},\hat{d}(c)).$$

Moreover, notice that $\hat{V}(c,.)$ is a particular solution to the family of nonhomegeneous ODEs parametrized by $c \in \mathbb{R}$:

$$[\mathcal{L}\phi(c,\cdot)](d) - g(c,d) \;\; = \;\; 0. \tag{4.6}$$

Therefore the general solution to (4.6) is in the form:

$$A(c)\psi(d) + B(c)\varphi(d) + \hat{V}(c,d),$$

for some real-valued constants $A(c)$, $B(c)$, which proves the structure of v in the continuation region, i.e. (4.4). □

Lemma 4.2. *The following properties hold:*

(i) $\lim_{d \downarrow d_{min}}(v(c,d) - \hat{V}(c,d)) = 0$*, for all* $c > \underline{c}$.

(ii) If $\bar{c} < \infty$*, then* $v(c,d) = \hat{V}(c,d)$ *for all* $c > \bar{c}$ *and* $d \in \mathcal{O}$.

(iii) If $\bar{c} = \infty$*, then* $\lim_{c \uparrow \infty}(v(c,d) - \hat{V}(c,d)) = 0$*, for all* $d \in \mathcal{O}$.

Proof. (i) Fix $c > \underline{c}$. By Lemma 4.1 $v(c,\cdot) \in C^2((d_{min}, \hat{d}(c)); \mathbb{R})$, and satisfies in a classical sense

$$[\mathcal{L}v(c,\cdot)](d) - g(c,d) \;\; = \;\; 0, \qquad \forall d \in (d_{min}, \hat{d}(c)). \tag{4.7}$$

Let $d \in (d_{min}, \hat{d}(c))$ and consider the stopping time $\tau_d = \inf\{t \geq 0 \mid D_t^d \geq \hat{d}(c)\}$. Since d_{min} is a not-entrance boundary for the diffusion D, we have (see e.g. [7, Ch II, Sec. 10]):

$$\tau_d \nearrow \infty \quad \text{a.s.} \qquad \text{when} \quad d \downarrow d_{min}. \tag{4.8}$$

By (4.7) and definition of τ_d, Itô's formula applied to $v(c, D_t^d)$ in the interval $[0, \tau_d)$ yields

$$v(c,d) \;\; = \;\; \mathbb{E}\Big[\int_0^{\tau_d} e^{-\rho t} g(c, D_t^d)\mathrm{d}t + e^{-\rho\tau_d} v(c, \hat{d}(c))\Big] \; \geq \; \mathbb{E}\Big[\int_0^{\tau_d} e^{-\rho t} g(c, D_t^d)\mathrm{d}t\Big].$$

Subtracting $\hat{V}(c,d)$ in both sides of the inequality above, we get

$$v(c,d) - \hat{V}(c,d) \;\; \geq \;\; \mathbb{E}\Big[\int_{\tau_d}^{\infty} e^{-\rho t} g(c, D_t^d)\mathrm{d}t\Big]$$

Taking the liminf for $d \downarrow d_{min}$, and using (4.8), we obtain

$$\liminf_{d \downarrow d_{min}} (v(c,d) - \hat{V}(c,d)) \;\; \geq \;\; 0,$$

and so the required limiting result since we always have $v \leq \hat{V}$ (see (3.2)).
(ii) Fix $c > \bar{c}$. Recalling (3.9), we then see that (c,d) lies in $\mathcal{C}$ for all $d \in \mathcal{O}$. Thus, $v(c,.)$ satisfies in a classical sense:

$$[\mathcal{L}v(c,\cdot)](d) - g(c,d) \;\; = \;\; 0, \qquad \forall d \in \mathcal{O}.$$

Thus, by applying Ito's formula to $v(c, D_t^d)$ between $t = 0$ and $T > 0$, we have:

$$v(c,d) \quad = \quad \mathbb{E}\Big[\int_0^T e^{-\rho t} g(c, D_t^d)\mathrm{d}t\Big] + \mathbb{E}\big[e^{-\rho T} v(T, D_T^d)\big]. \tag{4.9}$$

Now, from (3.3)-(3.4)-(3.5), we easily see that $\mathbb{E}\big[e^{-\rho T} v(T, D_T^d)\big]$ converges to zero as T goes to infinity. Therefore, by sending T to infinity into (4.9), and using monotone convergence theorem, we deduce that $v(c,d) = \hat{V}(c,d)$.

(iii) Suppose that $\bar{c} = \infty$. By Proposition 3.4-(1), we have $\hat{d}(c) \nearrow d_{max}$ when $c \nearrow \infty$. Let us consider the stopping time $\tau_c = \inf\{t \geq 0 \mid D_t^d \geq \hat{d}(c)\}$, which satisfies $\tau_c \nearrow \infty$ a.s. when $c \nearrow \infty$, since d_{max} is not-exit (see e.g. [7, Ch. II, Sec. 10]). Then, by arguing as in the proof of item (i) (with τ_c in place of τ_d), we obtain the desired claim. ☐

We can now provide the complete structure of the value function. Let us define

$$\underline{d} \quad := \quad \inf\{d \in \mathcal{O} \mid \hat{c}(d) > -\infty\}.$$

Theorem 4.1. *(Structure and properties of the value function)*
There exist functions $A : (\underline{c}, \infty) \to \mathbb{R}$, $z : (\underline{d}, d_{max}) \to \mathbb{R}$, such that

$$v(c,d) \quad = \quad \begin{cases} A(c)\psi(d) + \hat{V}(c,d) & \text{on } \bar{\mathcal{C}}, \\ -q_0 c + z(d), & \text{on } \mathcal{A}, \end{cases} \tag{4.10}$$

Moreover:

(i) The function A lies in $C^1((\underline{c}, \infty); \mathbb{R})$, and satisfies

$$\begin{cases} A(c) < 0, & c \in (\underline{c}, \bar{c}), \\ A(c) = 0, & c \geq \bar{c}, \end{cases} \tag{4.11}$$

(with the convention $\lim_{c\to\infty} A(c) = 0$ in the case $\bar{c} = \infty$). Furthermore, if there exists $d \in \mathcal{O}$ such that $\hat{c}(d) = \underline{c}$, then A can be extended to a function $C^1([\underline{c}, \infty); \mathbb{R})$.

(ii) z can be written in terms of v and $\hat{c}$ as

$$z(d) \quad = \quad v(\hat{c}(d), d) + q_0 \hat{c}(d). \tag{4.12}$$

(iii) The function $\hat{c}$ is right-continuous on $(\underline{d}, d_{max})$ and $\mathcal{C}$ is open.

Proof. *Structure of v in $\bar{\mathcal{C}}$.* By Lemma 4.1 and (3.18), we already know that there exist functions $A, B : (\underline{c}, \infty) \to \mathbb{R}$ such that

$$v(c,d) \quad = \quad A(c)\psi(d) + B(c)\varphi(d) + \hat{V}(c,d), \quad \text{in } \overset{\circ}{\mathcal{C}}.$$

Moreover, from Lemma 4.2 (i) and (4.3), we must have $B(c) = 0$. This proves the structure

$$v(c,d) \quad = \quad A(c)\psi(d) + \hat{V}(c,d), \quad \text{in } \overset{\circ}{\mathcal{C}}. \tag{4.13}$$

Let us now check that such structure also holds at the points of the boundary $\partial\mathcal{C}$.[4] Since $\hat{c}$ is nondecreasing, if $(c,d) \in \partial\mathcal{C}$, then one of the following case should hold:

---

[4] $\partial\mathcal{C}$ is intended as the boundary of $\mathcal{C}$ in $\mathcal{S}$.

(a) $d = \hat{d}(c) < d_{max}$,

(b) $c = \hat{c}(d)$ and $\{(c,d) \mid c > \hat{c}(d)\} \subset \mathring{\mathcal{C}}$,

(c) $\hat{d}(c') = \hat{d}(c)$ for $c' \in (c, c+\varepsilon)$ for some $\varepsilon > 0$.

In the case (a) the form (4.13) holds by Lemma 4.1. In the case (b) the structure (4.13) holds by continuity of v with respect to c and by the already proved structure in $\mathring{\mathcal{C}}$. In the case (c) the structure (4.13) holds by case (a) and by continuity of v with respect to c. In any case, we have proved that

$$v(c,d) \quad = \quad A(c)\psi(d) + \hat{V}(c,d), \qquad \text{in } \bar{\mathcal{C}}. \tag{4.14}$$

Structure of v in $\mathcal{A}$. This follows directly from the definition (3.8) of $\mathcal{A}$.

Let us now prove the remaining properties.

(i) The property that $A \in C^1((\underline{c},\infty);\mathbb{R})$ is consequence of the fact that $\hat{V}(\cdot,d)$ as well as $v(\cdot,d)$ (see Proposition 3.1) lie in $C^1((\hat{c}(d),\infty);\mathbb{R})$, together with the fact that $\psi(d) > 0$ for $d \in \mathcal{O}$. Let us prove (4.11). First of all, we see that $A \le 0$, since $v \le \hat{V}$ and $\psi > 0$. The fact that $A(c) = 0$, for $c \ge \bar{c}$ follows from Lemma 4.2 (ii). We prove now that $A(c) < 0$ for $c \in (\underline{c}, \bar{c})$. Fix some $c \in (\underline{c}, \bar{c})$. One can find $d_0 < d_{max}$ such that $c < \hat{c}(d_0)$. Since $\hat{c} \le \hat{c}_g$ and $g(\cdot, d_0)$ is strictly convex there exist $\varepsilon, \delta > 0$ such that

$$g_c(\xi, d_0) \quad \le \quad -\rho q_0 - \delta, \quad \forall \xi \in [c, c+\varepsilon].$$

Together with Assumption 2.2-(ii), we then get

$$g_c(\xi, d) \quad \le \quad -\rho q_0 - \delta, \quad \forall \xi \in [c, c+\varepsilon], \forall d > d_0,$$

and so

$$g(c+\varepsilon, d) \quad \le \quad g(c,d) + (-\rho q_0 - \delta)\varepsilon, \quad \forall d > d_0. \tag{4.15}$$

Let $d > d_0$ and consider the stopping time $\tau_d := \inf\{t \ge 0 \mid D_t^d \le d_0\}$. Since d_{max} is not-exit, we know that $\tau_d \nearrow \infty$ almost surely when d goes to d_{max} (see [7, Ch. II, Sec. 10]). Assume now on the contrary that $A(c) = 0$, so that $v(c,d) = \hat{V}(c,d)$ for all $d \in \mathcal{O}$. Take the control $I_t^\varepsilon \equiv \varepsilon$, so that

$$v(c,d) \quad \le \quad \mathbb{E}\Big[\int_0^{\tau_d} e^{-\rho t} g(c+\varepsilon, D_t^d)\mathrm{d}t + q_0\varepsilon + e^{-\rho\tau_d} v(c+\varepsilon, d_0)\Big].$$

Combining with (4.15), we then get

$$v(c,d) \quad \le \quad \mathbb{E}\Big[\int_0^{\tau_d} e^{-\rho t} g(c, D_t^d)\mathrm{d}t + (-\rho q_0 - \delta)\varepsilon \frac{1 - e^{-\rho\tau^d}}{\rho} + q_0\varepsilon + e^{-\rho\tau_d} v(c+\varepsilon, d_0)\Big].$$

Taking the limit for $d \uparrow d_{max}$, we get the required contradiction: $v(c,d) \le \hat{V}(c,d) - \varepsilon\delta/\rho$.

On the other hand, suppose that there exists $d \in \mathcal{O}$ such that $\hat{c}(d) = \underline{c}$. Then by continuity of v_c with respect to c we have

$$\lim_{c \downarrow \underline{c}} \left(A'(c)\psi(d) + \hat{V}_c(c,d)\right) \quad = \quad -q_0.$$

Since $\psi(d) > 0$, this implies that there exists a finite $\lim_{c \downarrow \underline{c}} A'(c)$, and therefore A can be extended to a function $C^1([\underline{c}, \infty); \mathbb{R})$.

(ii) Relation (4.12) follows from continuity of $v(\cdot, d)$ (Proposition 3.1), using (4.10) and evaluating v at $(\hat{c}(d), d) \in \mathcal{A}$.

(iii) Let $d_0 > \underline{d}$ and let $(d_n)_{n \geq 1}$ be a sequence s.t. $d_n \downarrow d_0$. We denote by $c_0 = \lim_{n \to \infty} \hat{c}(d_n)$. This limit exists by monotonicity of $\hat{c}$ and is larger than $\hat{c}(d_0)$. Hence, we only need to check that c_0 is smaller than $\hat{c}(d_0)$. By definition of $\mathcal{A}$ and $\hat{c}(d_n)$, the point $(\hat{c}(d_n), d_n)$ lies on $\mathcal{A}$, so

$$v_c(\hat{c}(d_n), d_n) = -q_0, \quad \forall n \geq 1. \tag{4.16}$$

Moreover, by (3.10), $(\hat{c}(d_n), d_n)$ also lies in $\bar{\mathcal{C}}$, so

$$(c_0, d_0) \;=\; \lim_{n \to \infty} (\hat{c}(d_n), d_n) \;\in\; \bar{\mathcal{C}}. \tag{4.17}$$

Now, as consequence of Lemma 3.1 and (4.14), the function v_c is continuous on $\bar{\mathcal{C}}$. Therefore, by (4.16) and (4.17), we have:

$$v_c(c_0, d_0) \;=\; -q_0,$$

which proves that $(c_0, d_0) \in \mathcal{A}$, i.e. $\hat{c}(d_0) \geq c_0$, and so the right-continuity of $\hat{c}$. Finally, the fact that $\mathcal{C}$ is open follows from right-continuity and monotonicity of $\hat{c}$. $\square$

The second main result provides the existence and an explicit description of the optimal irreversible investment.

Theorem 4.2. *(Optimal control)*
Let $(c, d) \in \mathcal{S}$, and consider the process $I^* \in \mathcal{I}$ defined by

$$I_t^* = \left[\hat{c}\left(\sup_{0 \leq s \leq t} D_s^d \right) - c \right]^+ .$$

Then I^* is an optimal control for the value function at (c, d).

Proof. Fix $(c, d) \in \mathcal{S}$, and denote by $C^* = C^{c, I^*}$ the capacity process starting from c and controlled by $I^* \in \mathcal{I}$. We claim that

(a) $(C_t^*, D_t^d) \in \bar{\mathcal{C}}$, a.s. for all $t \geq 0$,

(b) $\int_0^\infty e^{-\rho t} \mathbf{1}_{\{(C_t^*, D_t^d) \in \mathcal{C}\}} \mathrm{d}I_t^* = 0$, a.s.

Indeed, by monotonicity of the function $\hat{c}$, we have

$$C_t^* \;=\; c + \Big[\hat{c}\big(\sup_{0 \leq s \leq t} D_s^d \big) - c \Big]^+ \;\geq\; \hat{c}(D_t^d),$$

which shows (a) by (3.10). Let us now prove the assertion (b). Fix $\omega \in \Omega$ and suppose that $(C_t^*(\omega), D_t^d(\omega)) \in \mathcal{C}$, i.e. $C_t^*(\omega) > \hat{c}(D_t^d(\omega))$ by (3.9). We distinguish two cases:

(i) $\hat{c}\big(\sup_{0\leq s\leq t} D^d_s(\omega)\big) \geq c$. In this case we have

$$\hat{c}\Big(\sup_{0\leq s\leq t} D^d_s(\omega)\Big) \;=\; C^*_t(\omega) \;>\; \hat{c}(D^d_t(\omega)),$$

and by monotonicity of $\hat{c}$ this means that

$$\sup_{0\leq s\leq t} D^d_s(\omega) \;>\; D^d_t(\omega).$$

Then, by continuity of $D^d_\cdot(\omega)$, we deduce that $r \mapsto \sup_{0\leq s\leq r} D^d_s(\omega)$ is constant in the interval $[t, t+\varepsilon(\omega))$ for some $\varepsilon(\omega) > 0$, and the same holds true for $r \mapsto I^*_r(\omega)$ by definition of I^*.

(ii) $\hat{c}\left(\sup_{0\leq s\leq t} D^d_s(\omega)\right) < c$. In this case we have

$$c \;=\; C^*_t(\omega) \;>\; \hat{c}(D^d_t(\omega)).$$

Hence, by monotonicity and right-continuity of $\hat{c}$, and by continuity of $D^d_\cdot(\omega)$, there exists some $\varepsilon(\omega) > 0$ s.t.

$$c \;>\; \sup_{t\leq r\leq t+\varepsilon(\omega)} \hat{c}(D^d_r(\omega)).$$

Together with the assumption of this case, this yields $c > \sup_{0\leq r\leq t+\varepsilon(\omega)} \hat{c}(D^d_r(\omega))$, and so: $I^*_r(\omega) = 0$ for all $r \in [0, t+\varepsilon(\omega))$.

In any case, we have shown that, if $(C^*_t, D^d_t) \in \mathcal{C}$, then I^* is constant in a right (stochastic) neighborhood of t, which shows (b). Now let $\mathcal{N}$ be the (at most countable) set of discontinuity points of $\hat{c}$. Due to continuity of $\sup_{0\leq s\leq t} D^d_s$ and by definition of I^*,

(c) the jumps $\Delta I^*_t = \Delta C^*_t$ of I^*_t and C^*_t occur only in the following cases:

- either at time 0 when $c < \hat{c}(d)$, and in this case $\Delta I^*_0 = \hat{c}(d) - c$;
- or when $D^d_t \in \mathcal{N}$ and $C^{c,*}_{t^-} < \hat{c}(D^d_t)$, and in this case $\Delta I^*_t = \hat{c}(D^d_t) - C^*_{t^-}$.

Let us now prove that

$$v(c,d) \;\;\geq\;\; \mathbb{E}\Big[\int_0^\infty e^{-\rho t}\Big(g(C^*_t, D^d_t) + q_0 \mathrm{d}I^*_t\Big)\Big]. \tag{4.18}$$

Consider the (bounded) stopping time $\tau_n = \inf\{t \geq 0 \mid C^*_t \wedge D^d_t \geq n\} \wedge n$, and notice that $\tau_n \nearrow \infty$ a.s. when n goes to infinity. From (4.14), we see that $v \in C^{1,2}(\bar{\mathcal{C}};\mathbb{R})$. Thus, by the assertion (a), we may apply Itô's formula (see Theorem 32, pag. 79, in [27]) to $e^{-\rho t}v(C^*_t, D^d_t)$ between 0 and τ_n, take expectation, and obtain (after observing that the stochastic integral over the interval $[0, \tau_n \wedge T)$ vanishes in expectation):

$$\begin{aligned} v(c,d) \;\;=\;\; & \mathbb{E}\Big[e^{-\rho\tau_n} v(C^*_{\tau_n\wedge T}, D^d_{\tau_n\wedge T})\Big] \;+\; \mathbb{E}\Big[\int_0^{\tau_n} e^{-\rho t}[\mathcal{L}v(C^*_t,\cdot)](D^d_t)\mathrm{d}t\Big] \\ & - \mathbb{E}\Big[\int_0^{\tau_n} e^{-\rho t} v_c(C^*_t, D^d_t)\mathrm{d}I^*_t\Big] \\ & - \mathbb{E}\Big[\sum_{0\leq t\leq\tau_n} e^{-\rho t}(v(C^*_t, D^d_t) - v(C^*_{t^-}, D^d_t) - v_c(C^*_t, D^d_t)\Delta C^*_t)\Big], \end{aligned} \tag{4.19}$$

Now observe that $[\mathcal{L}v(c',\cdot)](d') = g(c',d')$ for (c',d') in $\mathcal{C}$ but also in $\bar{\mathcal{C}}$ by continuity of g and (4.14). Together with (a) this implies that

$$\mathbb{E}\Big[\int_0^{\tau_n} e^{-\rho t}[\mathcal{L}v(C_t^*,\cdot)](D_t^d)\mathrm{d}t\Big] \;\;=\;\; \mathbb{E}\Big[\int_0^{\tau_n} e^{-\rho t}\big(g(C_t^*,D_t^d)\mathrm{d}t\Big].$$

By (b) and since $v_c = -q_0$ in $\mathcal{A}$, we have

$$\mathbb{E}\Big[\int_0^{\tau_n} e^{-\rho t}v_c(C_t^*,D_t^d)\mathrm{d}I_t^*\Big] \;\;=\;\; -\,\mathbb{E}\Big[\int_0^{\tau_n} e^{-\rho t}q_0\mathrm{d}I_t^*\Big].$$

Moreover, by (c) we have

$$v(C_t^*,D_t^d) - v(C_{t-}^*,D_t^d) - v_c(C_t^*,D_t^d)\Delta C_t^* \;\;=\;\; 0, \quad a.s.,\ \forall t \geq 0.$$

Therefore by nonnegativity of v and (4.19), we have

$$v(c,d) \;\;\geq\;\; \mathbb{E}\Big[\int_0^{\tau_n} e^{-\rho t}\big(g(C_t^*,D_t^d)\mathrm{d}t + q_0\mathrm{d}I_t^*\big)\Big].$$

By sending n to infinity, and from monotone convergence, we get the inequality (4.18). Since the opposite inequality always hold by definition of v, this proves the equality, i.e. that I^* is an optimal control. □

The picture below represents a possible shape of the solution. The state space region $\mathcal{S}$ is the half-plane on the right of the vertical dotted line. When the system lies in the continuation region $\mathcal{C}$, it moves along the horizontal lines and no action is taken. Whenever the system touches the boundary $\partial\mathcal{C}$, the optimal control (acting along the vertical lines as indicated by the arrows in the picture) consists in investing the minimal amount of money to keep the system in $\bar{\mathcal{C}}$.



5 The quadratic case: second-order smooth fit and explicit solution

Theorem 4.1 and continuity of v, v_c with respect to c yield a couple of optimality conditions. Indeed, the functions $A, \hat{c}, z$ must satisfy for every $d \in (\underline{d}, d_{max})$:

$$\begin{cases} \textbf{(i)} \;\; A(\hat{c}(d))\psi(d) + \hat{V}(\hat{c}(d), d) \;=\; -q_0\hat{c}(d) + z(d), \\ \\ \textbf{(ii)} \;\; A'(\hat{c}(d))\psi(d) + \hat{V}_c(\hat{c}(d), d) \;=\; -q_0. \end{cases} \tag{5.1}$$

It is clear that one cannot expect that the two conditions above provide a way either to find the value function or just the optimal boundary $\hat{c}$, as they relate three functions by two equations. A third condition is needed and should be derived from some other suitable smoothness property of the value function at the optimal boundary. To this end, we notice by Theorem 4.1 that

$$\frac{\partial}{\partial d} v_c(c, d) = 0 \;\text{ in }\; \overset{\circ}{\mathcal{A}}\,. \tag{5.2}$$

Therefore, a requirement of a smooth fit condition of the second order mixed derivative of v at the optimal boundary would imply

$$\lim_{c \downarrow \hat{c}(d)} v_{cd}(c, d) \;=\; 0, \qquad \forall d \in (\underline{d}, d_{max}). \tag{5.3}$$

(Remark 5.1 below makes this point more precise).

The purpose of the present section is indeed to prove (5.3). However, we need to further specify our assumptions, restricting to the quadratic cost case:

$$g(c, d) \;=\; \frac{1}{2}(c^2 - 2\beta_0(d)c + \alpha_0(d)), \tag{5.4}$$

where α_0, β_0 are continuous functions. Within this section we assume that g has the structure (5.4) and we do not repeat this assumption in the statements of the results. We assume that the function β_0 is nondecreasing, so that Assumption 2.2-(ii) holds true, and we denote by

$$\alpha(d) \;:=\; \mathbb{E}\Big[\int_0^\infty e^{-\rho t}\alpha_0(D_t^d)\mathrm{d}t\Big], \qquad \beta(d) \;:=\; \mathbb{E}\Big[\int_0^\infty e^{-\rho t}\beta_0(D_t^d)\mathrm{d}t\Big], \tag{5.5}$$

and notice that $\alpha, \beta \in C^2((d_{min}, d_{max}); \mathbb{R})$ as the diffusion D is nondegenerate. The function $\hat{V}$ is written in this case as:

$$\hat{V}(c, d) \;=\; \frac{1}{2}\Big(\frac{1}{\rho}c^2 - 2\beta(d)c + \alpha(d)\Big). \tag{5.6}$$

Given a function $w \in C(\mathbb{R}; \mathbb{R})$, let us denote by

$$[\Delta^2 w](x; \varepsilon) \;=\; \frac{1}{\varepsilon^2}[w(x+\varepsilon) + w(x-\varepsilon) - 2w(x)], \quad x \in \mathbb{R},\; \varepsilon > 0.$$

The following Lemma, which relies on assumption (5.4), enables us to obtain further regularity of the value function with respect to c (Corollary 5.1), which is crucial to prove then (5.3).

Lemma 5.1. *We have*

$$0 \;\le\; [\Delta^2 v(\cdot,d)](c;\varepsilon) \;\;\le\;\; \frac{1}{\rho}, \qquad \forall c \in \mathbb{R},\ \forall \varepsilon > 0,\ \forall d \in \mathcal{O}.$$

Proof. The estimate from below is a straightforward consequence of the convexity of v with respect to c. Let us prove the estimate from above. Let $(c,d) \in \mathcal{S}$, $\varepsilon > 0$ and $I \in \mathcal{I}$. By using the fact that $g_{cc} \equiv 1$ under (5.4), we have

$$\begin{aligned} & \frac{1}{\varepsilon^2}\left[G(c+\varepsilon,d;I) + G(c-\varepsilon,d;I) - 2G(c,d;I)\right] \qquad (5.7)\\ = \;\; & \mathbb{E}\Big[\int_0^\infty e^{-\rho t}\Big[\frac{1}{\varepsilon^2}\big(g(C_t^{c+\varepsilon,I},D_t^d) + g(C_t^{c-\varepsilon,I},D_t^d) - 2g(C_t^{c,I},D_t^d)\big)\Big)\mathrm{d}t\Big] \;=\; \frac{1}{\rho}. \end{aligned}$$

Since

$$v(c+\varepsilon,d) + v(c-\varepsilon,d) - 2G(c,d;I) \;\;\le\;\; G(c+\varepsilon,d;I) + G(c-\varepsilon,d;I) - 2G(c,d;I),$$

we get from (5.7):

$$\frac{1}{\varepsilon^2}\left[v(c+\varepsilon,d) + v(c-\varepsilon,d) - 2G(c,d;I)\right] \;\;\le\;\; \frac{1}{\rho}, \quad \forall c \in \mathbb{R},\ \forall d \in \mathcal{O},\ \forall \varepsilon > 0.$$

Taking the supremum over $I \in \mathcal{I}$, this proves the required upper-estimate. □

Lemma 5.1 implies that $v_c(\cdot,d)$ is Lipschitz continuous for each $d \in \mathcal{O}$. Together with Theorem 4.1, we immediately get the following regularity result.

Corollary 5.1. *The derivative function $A' : (\underline{c},\infty) \to \mathbb{R}$, where A is the function defined in Theorem 4.1, is locally Lipschitz. In other terms $A \in W^{2,\infty}_{loc}((\underline{c},\infty);\mathbb{R})$.*

We are now able to prove the second order smooth-fit result on the value function.

Proposition 5.1. *The relation* (5.3) *holds true:*

$$\lim_{c \downarrow \hat{c}(d)} v_{cd}(c,d) \;\;=\;\; 0, \qquad \forall d \in (\underline{d}, d_{max}).$$

Proof. Let us consider the function w defined on $(\underline{c},\infty) \times \mathcal{O}$ by

$$w(c,d) \;\;:=\;\; A(c)\psi(d) + \hat{V}(c,d).$$

We have $w = v$ over $\bar{\mathcal{C}}$, so the claim is equivalent to prove that

$$\lim_{c \downarrow \hat{c}(d)} w_{cd}(c,d) \;\;=\;\; 0, \qquad \forall d \in \mathcal{O}.$$

We notice that the limit above always exists and coincides with $w_{cd}(\hat{c}(d),d)$ due to continuity of w_{cd} (see Theorem 4.1). Suppose by contradiction that there exists $d_0 \in (\underline{d}, d_{max})$ such that the limit above is different from 0, in which case, by Proposition 3.2-(ii), we should have

$$w_{cd}(\hat{c}(d_0),d_0) \;\;<\;\; 0. \qquad (5.8)$$

Due to right continuity of $\hat{c}$ (Proposition 4.1-(iii)), we may find $d_1 > d_0$ such that $\hat{c}$ is continuous in $[d_0, d_1]$. Under (5.8) and by continuity of w_{cd}, we may find $\varepsilon > 0$ and $d_2 \in (d_0, d_1)$ such that, denoting by $c_0 = \hat{c}(d_0)$, $c_2 = \hat{c}(d_2)$, we have

$$w_{cd}(c,d) \quad \leq \quad -\varepsilon, \quad \forall (c,d) \in [c_0, c_2] \times [d_0, d_2].$$

Consider now the function $\tilde{w}(c,d) = w(c,d) + q_0 c$, and observe that we have as well

$$\tilde{w}_{cd}(c,d) \quad \leq \quad -\varepsilon, \quad \forall (c,d) \in [c_0, c_2] \times [d_0, d_2]. \tag{5.9}$$

Moreover, by definition of $\hat{c}$, we have

$$\tilde{w}_c(\hat{c}(d), d) \quad = \quad 0, \quad \forall d \in [d_0, d_2]. \tag{5.10}$$

We then have the following facts:

1. From (5.9), the function $\hat{c}$ is strictly increasing in the interval $[d_0, d_2]$ and therefore can be inverted. The inverse function $\hat{d} = \hat{c}^{-1} : [c_0, c_2] \to [d_0, d_2]$ is continuous and strictly increasing as well.
2. By (5.10), and Corollary 5.1, we can apply a (generalized) implicit function theorem to $\tilde{w}_c$ and get in a generalized sense

   $$\hat{d} \in W^{1,\infty}([c_0, c_2]; [d_0, d_2]), \qquad \hat{d}'(c) \;=\; -\frac{\tilde{w}_{cc}(c, \hat{d}(c))}{\tilde{w}_{cd}(c, \hat{d}(c))},$$

   with

   $$\|\hat{d}'\|_\infty \;\leq\; \frac{\mathrm{esssup}_{c \in [c_0, c_2]} \tilde{w}_{cc}(c, \hat{d}(c))}{\varepsilon} \quad =: \quad M_\varepsilon \;<\; \infty.$$

3. By item 2, $\hat{d}$ is absolutely continuous. Hence, by using also item 1, we see that the set

   $$\mathcal{N} \quad := \quad \{c \in (c_0, c_2) \mid \exists\, \hat{d}'(c) \in (0, M_\varepsilon]\}$$

   has full Lebesgue measure in $[c_0, c_2]$.
4. Denoting by

   $$\hat{d}(\mathcal{N}) \quad = \quad \{d = \hat{d}(c) \mid c \in \mathcal{N}\},$$

   we see by the above items that

   $$\begin{cases} \text{(i) } \hat{d}(\mathcal{N}) \text{ is dense in } [d_0, d_2], \\ \text{(ii) } \hat{c} \text{ is differentiable at each } d \in \hat{d}(\mathcal{N}), \\ \text{(iii) } \hat{c}'(d) \in [1/M_\varepsilon, \infty), \ \forall d \in \hat{d}(\mathcal{N}). \end{cases} \tag{5.11}$$

Let us now consider the function $d \in [d_0, d_2] \mapsto v(c_0, d)$. Since $\hat{c}$ is nondecreasing, the segment $\{(c_0, d) \mid d \in [d_0, d_2]\}$ is contained in $\mathcal{A}$. Hence, Theorem 4.1 yields

$$v(c_0, d) = -q_0 c_0 + z(d). \tag{5.12}$$

Apply the chain rule at the points of $\hat{d}(\mathcal{N})$ to

$$[d_0, d_2] \to \mathbb{R}, \qquad d \mapsto z(d) \;=\; v(\hat{c}(d), d) + q_0 \hat{c}(d) \;=\; w(\hat{c}(d), d) + q_0 \hat{c}(d).$$

Then, the function z is differentiable at the points of $\hat{d}(\mathcal{N})$ and

$$z'(d) \quad = \quad w_c(\hat{c}(d), d)\hat{c}'(d) + w_d(\hat{c}(d), d) + q_0 \hat{c}'(d), \quad \forall d \in \hat{d}(\mathcal{N}).$$

By definition of $\hat{c}$, we have: $w_c(\hat{c}(d), d) = v_c(\hat{c}(d), d) = -q_0$ for every $d \in \mathcal{O}$, and so

$$z'(d) \;=\; v_d(\hat{c}(d), d), \quad \forall d \in \hat{d}(\mathcal{N}).$$

Together with (5.12), this shows the existence of $v_d(c_0, d)$ for each $d \in \hat{d}(\mathcal{N})$ and the equality

$$v_d(c_0, d) \;=\; z'(d) \;=\; w_d(\hat{c}(d), d), \quad \forall d \in \hat{d}(\mathcal{N}). \tag{5.13}$$

On the other hand, by using the same argument for the use of the chain rule, we can get from (5.13) the existence of $v_{dd}(c_0, d)$ for each $d \in \hat{d}(\mathcal{N})$ and the equality

$$v_{dd}(c_0, d) \;=\; z''(d) \;=\; w_{dd}(\hat{c}(d), d) + w_{cd}(\hat{c}(d), d)\, \hat{c}'(d), \quad \forall d \in \hat{d}(\mathcal{N}).$$

Therefore, from (5.9), we get

$$v_{dd}(c_0, d) \;\leq\; w_{dd}(\hat{c}(d), d) - \varepsilon / M_\varepsilon, \quad \forall d \in \hat{d}(\mathcal{N}). \tag{5.14}$$

Now the viscosity subsolution property of v and (5.12)-(5.13)-(5.14) yield

$$\rho v(c_0, d) - \mu(d) w_d(\hat{c}(d), d) - \frac{1}{2}\sigma(d)^2 [w_{dd}(\hat{c}(d), d) - \varepsilon / M_\varepsilon] \;\leq\; g(c_0, d), \quad \forall d \in \hat{d}(\mathcal{N}). \tag{5.15}$$

Taking a sequence $(\alpha_n) \subset \hat{d}(\mathcal{N})$ such that $\alpha_n \downarrow d_0$ (this can be done by (5.11)-(i)) and passing to the limit in (5.15) evaluated at $d = \alpha_n$ we obtain:

$$\rho v(c_0, d_0) - \mu(d_0) w_d(c_0, d_0) - \frac{1}{2}\sigma(d_0)^2 [w_{dd}(c_0, d_0) - \varepsilon / M_\varepsilon] \;\leq\; g(c_0, d_0). \tag{5.16}$$

On the other hand, recall that $\mathcal{L}w = g$ on $\mathcal{C}$, as $v = w$ therein. Therefore, since $w \in C^{1,2}(\mathcal{S}; \mathbb{R})$ and since $(c_0, d_0) \in \partial\mathcal{C}$, by continuity of g we must also have

$$\rho v(c_0, d_0) - \mu(d_0) w_d(c_0, d_0) - \frac{1}{2}\sigma(d_0)^2 w_{dd}(c_0, d_0) \quad = \quad g(c_0, d_0),$$

which is in contradiction with (5.16) as $\sigma^2(d_0) > 0$, and the claim is proved. □

Proposition 5.1 can be used to add a third condition to (5.1): indeed, by (4.14), the relation (5.3) yields

$$A'(\hat{c}(d))\psi'(d) + \hat{V}_{cd}(\hat{c}(d), d) \quad = \quad 0, \quad \forall d \in (\underline{d}, d_{max}),$$

so that the functions $A, \hat{c}, z$ must satisfy for all $d \in (\underline{d}, d_{max})$

$$\begin{cases} \textbf{(i)} \;\; A(\hat{c}(d))\psi(d) + \hat{V}(\hat{c}(d), d) \; = \; -q_0\hat{c}(d) + z(d), \\ \\ \textbf{(ii)} \;\; A'(\hat{c}(d))\psi(d) + \hat{V}_c(\hat{c}(d), d) \; = \; -q_0, \\ \\ \textbf{(iii)} \;\; A'(\hat{c}(d))\psi'(d) + \hat{V}_{cd}(\hat{c}(d), d) \; = \; 0. \end{cases} \tag{5.17}$$

This is a set of necessary optimality conditions, which are sufficient for identifying the value function.

Theorem 5.1. *The functions $\hat{c}, A, z$ of Theorem 4.1 are determined as follows:*

(a) The optimal boundary function $\hat{c}$ is explicitly given by

$$\hat{c}(d) \; = \; \rho \left[\beta(d) - \frac{\psi(d)}{\psi'(d)}\beta'(d) - q_0 e^{\rho h} \right], \quad d \in \mathcal{O}. \tag{5.18}$$

(b) The function A is given by[5]

$$A(c) \; = \; -\int_c^{\bar{c}} \frac{\beta'(\hat{d}(\xi))}{\psi'(\hat{d}(\xi))} \mathrm{d}\xi, \qquad c \in (\underline{c}, \bar{c}), \tag{5.19}$$

where $\hat{d}$ is the pseudo-inverse of $\hat{c}$.

(c) The function z is equal to

$$z(d) \; = \; A(\hat{c}(d))\psi(d) + \hat{V}(\hat{c}(d), d) + q_0\hat{c}(d). \tag{5.20}$$

Proof. (a) Multiplying (5.17)-(iii) by ψ/ψ' and subtracting to (5.17)-(ii), we get (5.18).
(b) From (5.17)-(iii), (4.11) and the expression (5.6) of $\hat{V}$, we get (5.19) by integration.
(c) The last equality (5.20) is derived from (a)-(b), and (5.1)-(i). □

Remark 5.1. 1. As consequence of Theorem 5.1, we see that the optimal boundary $\hat{c}$ is continuous (and actually smooth), and so

$$\overset{\circ}{\mathcal{A}} \;\; = \;\; \{(c, d) \in \mathcal{S} \mid c < \hat{c}(d)\}.$$

Then, Proposition 5.1 together with (5.2) yield

$$\lim_{c \downarrow \hat{c}(d)} v_{cd}(c, d) \;\; = \;\; \lim_{c \uparrow \hat{c}(d)} v_{cd}(c, d), \qquad \forall d \in (\underline{d}, d_{max}),$$

which shows clearly the smooth-fit of the mixed second derivative of the value function.

2. We observe that (5.17) also implies the continuity of v_d, v_{dd} at $\partial\mathcal{C}$. Indeed, differentiating (5.17)-(i) with respect to d and subtracting (5.17)-(ii) multiplied by $\hat{c}'(d)$ we get

$$A(\hat{c}(d))\psi'(d) + \hat{V}_d(\hat{c}(d), d) \; = \; z'(d),$$

---

[5] It can be eventually extended by continuity to $\underline{c}$ and $\bar{c}$, if they are finite.

which means the continuity of v_d at $\partial\mathcal{C}$. Then, differentiating (5.21) with respect to d and subtracting (5.17)-(iii) multiplied by $\hat{c}'(d)$, we have

$$A(\hat{c}(d))\psi''(d) + \hat{V}_{dd}(\hat{c}(d), d) \;=\; z''(d),$$

meaning the continuity of v_{dd} at $\partial\mathcal{C}$. □

Remark 5.2. We comment now about possible relaxing of assumption (5.4). A first natural question arising is if the smooth fit principle stated in Proposition 5.1 is true even without assuming that structure for g. As we have outlined at the beginning of this section, assumption (5.4) allows to obtain a good regularity of the value function with respect to c needed in the proof of Proposition 5.1. On the other hand, we are quite confident that it holds also in other cases, maybe the proof passing through other arguments.

Supposing to have a positive answer to this first question, a second natural question is if this smooth fit principle is still sufficient to uniquely individuate the value function together with the other optimality conditions. The optimality conditions (5.17) remain the same, so the question just corresponds to ask if uniqueness of solutions holds for the system. Arguing as in the proof of item (a) of Theorem 5.1, one has to deal, for given $d \in \mathcal{O}$ with the solvability with respect to c of

$$\frac{\psi(d)}{\psi'(d)}\hat{V}_{cd}(c, d) - \hat{V}_c(c, d) \;=\; q_0. \tag{5.21}$$

We already know that there is a solution to the equation above for each $d > \underline{d}$, as $\hat{c}(d)$ must solve this equation (since it is derived from *necessary* conditions of optimality). Therefore the question reduces to the problem of uniqueness of solutions to (5.21) (at least in the interval $(-\infty, \hat{c}_g(d)]$, as by Proposition 3.3 we know that $\hat{c}(d) \le \hat{c}_g(d)$; observe also that $\hat{c}_g(d)$ is in principle easily computable). Once one knows that this uniqueness holds true, then one can still argue as in the remaining proof of Theorem 5.1 to get the complete expression of the functions A, z individuating the value function. □

A basic example: the GBM case

We end this paper by a simple and explicit illustration of our Theorem 5.1 to the case when the demand is modeled as a geometric Brownian motion:

$$\mathrm{d}D_t \;=\; \mu D_t \mathrm{d}t + \sigma D_t \mathrm{d}W_t, \quad \mu \in \mathbb{R},\ \sigma > 0,$$

with initial datum $d > 0$. In this case $\mathcal{O} = (0, \infty)$. Moreover, assume that

$$g(c, d) \;=\; \frac{1}{2}(c - d)^2,$$

and, according to Assumption 3.4, assume that

$$\rho > 2\mu + \sigma^2. \tag{5.22}$$

Then $\hat{V}$ is the quadratic function equal to

$$\hat{V}(c,d) \;=\; \frac{1}{2}\Big(\frac{1}{\rho-2\mu-\sigma^2}d^2 - \frac{2}{\rho-\mu}dc + \frac{1}{\rho}c^2\Big).$$

The increasing fundamental solution to

$$[\mathcal{L}\phi](d) \;:=\; \rho\phi - \mu d\phi' - \frac{1}{2}\sigma^2 d^2\phi'' \;=\; 0,$$

is given by

$$\psi(d) \;=\; d^m$$

where m is the positive root of the equation $\rho - \mu m - \frac{1}{2}\sigma^2 m(m-1) = 0$, and explicitly given by

$$m \;=\; -\frac{\mu}{\sigma^2} + \frac{1}{2} + \sqrt{\Big(-\frac{\mu}{\sigma^2}+\frac{1}{2}\Big)^2 + \frac{2\rho}{\sigma^2}}$$

Notice that $m > 2$ by (5.22). From Theorems 4.1 and 5.1, the value function v has the explicit form

$$v(c,d) \;=\; \begin{cases} A(c)d^m + \hat{V}(c,d), & \text{on } c > \hat{c}(d), \\ -q_0 c + z(d), & \text{on } c \leq \hat{c}(d). \end{cases}$$

where the functions $A, \hat{c}, z$ are given by

$$\begin{aligned} \hat{c}(d) &= ad - b, \qquad d > 0, \\ A(c) &= -\frac{a^{m-1}}{m(m-m-2)}\frac{1}{\rho-\mu}(c+b)^{2-m}, \qquad c > -b, \\ z(d) &= A(ad-b)d^m + \hat{V}(ad-b,d) + q_0(\alpha d - b), \qquad d > 0, \end{aligned}$$

with

$$a \;=\; \frac{m-1}{m}\frac{\rho}{\rho-\mu}, \qquad b = \rho q_0.$$

Acknowledgement. The authors would like to thank René Aïd and Bertrand Villeneuve for useful discussions. Salvatore Federico also thanks Giorgio Ferrari for comments on the literature and for suggesting the possible extension of the model described in Remark 2.2.

References

[1] Abel A.B. and J. C. Eberly, *Optimal investment with costly reversibility*, Review of Economic Studies, Vol. 63, (1996), pp. 581–593.

[2] Aïd R., Federico S., Pham H. and B. Villeneuve, *Technology choices and investment with time to build*, work in progress.

[3] Alvarez L.H., *Irreversible capital accumulation under interest rate uncertainty*, Mathematical Methods of Operations Research, Vol. 72 (2009), No. 2,pp.249–271.

[4] Alvarez L.H., *Optimal capital accumulation under price uncertainty and costly reversibility*, Journal of Economics, Dynamics and Control, Vol. 35 (2011), No. 10, pp. 1769–1788.

[5] Baldursson F.M. and I. Karatzas, *Irreversible investment and industry equilibrium*, Finance and Stochastics, Vol. 1 (1997), No. 1, pp.69–89.

[6] Bank P., *Optimal control under a dynamic fuel constraint*, SIAM Journal on Control and Optimization, Vol. 44 (2005), No. 4, pp.1529–1541.

[7] Borodin A.N. and P. Salminen, *Handbook of Brownian motion - Facts and formulae*, Second edition (2002), Birkhäuser.

[8] Bouchard B. and N. Touzi, *Weak dynamic programming principle for viscosity solutions*, SIAM Journal on Control and Optimization, Vol. 49 (2011), No. 3, pp. 948–962.

[9] Cannarsa P. and C. Sinestrari, *Semiconcave functions, Hamilton-Jacobi equations and optimal control*, Progress in Nonlinear Differential Equations and their Application (2004), Birkhäuser.

[10] Chiarolla M.B. and U.G. Haussman, *Explicit solution of a stochastic irreversible investment problem and its moving threshold*, Mathematics of Operations Research, Vol. 30 (2005), No. 1, pp. 91–108.

[11] Chiarolla M.B. and U.G. Haussman, *On a stochastic irreversible investment problem*, SIAM Journal on Control and Optimization, Vol. 48 (2009), No. 2, 438–462.

[12] Chow P.L., Menaldi J.L. and M. Robin, *Additive control of stochastic linear systems with finite horizon*, SIAM Journal on Control and Optimization, Vol. 23 (1985), No. 6, pp. 858–899.

[13] Davis M.H., Dempster M.A.H., Sethi S.P. and D. Vermes, *Optimal capacity expansion under uncertainty*, Advances in Applied Probability, Vol. 19 (1987), pp. 156–176.

[14] Dixit A.K. and R.S. Pindyck, *Investment under uncertainty*, Princeton University Press (1994).

[15] Fleming W. H. and H. M. Soner, *Controlled Markov Processes and Viscosity Solutions* (2006), Springer-Verlag.

[16] Guo X. and H. Pham, *Optimal partially reversible investments with entry decision and general production function*, Stochastic Processes and their Applications, Vol. 115 (2005), No. 5, pp. 705-736.

[17] Guo X. and P. Tomecek, *A class of singular control problems and the smooth fit principle*, SIAM Journal on Control and Optimization, Vol. 47 (2008), No. 6, pp. 3076–3099.

[18] Guo X. and G. L. Wu, *Smooth fit principle for impulse control of multi-dimensional diffusion processes*, SIAM Journal on Control and Optimization, Vol. 48 (2009), No. 2, pp. 594-617.

[19] Haussman U.G. and W. Suo, *Singular optimal stochastic controls I: existence*, SIAM Journal on Control and Optimization, Vol. 33 (1995), No. 3, pp. 916-936.

[20] Haussman U.G. and W. Suo, *Singular optimal stochastic controls II: dynamic programming*, SIAM Journal on Control and Optimization, Vol. 33 (1995),Vol. 33, No. 3, pp. 937-959.

[21] Karatzas I., *A class of singular stochastic control problems*, Advances in Applied Probability, Vol. 15 (1983), No. 2, pp. 225–254.

[22] Løkka A. and M. Zervos, *A model for the long-term optimal capacity level of an investment project*, International Journal of Theoretical and Applied Finance, Vol. 14 (2011), No. 2,pp. 187–196.

[23] Manne A.S., *Capacity expansion and probabilistic growth*, Econometrica, Vol. 29 (1961), No. 4, pp. 632–649.

[24] Merhi A. and M. Zervos, *A model for reversible investment capacity expansion*, SIAM Journal on Control and Optimization, Vol. 46 (2007), No. 3, pp. 839–876.

[25] Øksendal A., *Irreversible investment problems*, Finance and Stochastics, Vol. 4 (2000), No. 2, pp. 223–250.

[26] Pham H., *Continuous-time stochastic control and applications with financial applications*, Series Stochastic Modeling and Applied Probability, Vol. 61 (2009), Springer.

[27] Protter P., *Stochastic integration and differential equations*, Second edition (2004), Springer.

[28] Riedel F. and X. Su, *On irreversible investment*, Finance and Stochastics, Vol. 15 (2011), No. 4, pp. 607–633.

[29] Shreve S. and H.M. Soner, *Regularity of the value function for a two-dimensional singular stochastic control problem*, SIAM J. Control and Optimization, Vol. 27 (1989), pp. 876–907.

[30] Shreve S. and H.M. Soner, *Optimal investment and consumption with transactions costs*, The Annals of Applied Probability, Vol. 4 (1994), No. 3, pp. 609–692.

[31] Wang H., *Capacity expansion with exponential jump diffusion process*, Stochastics and Stochastics Reports, Vol. 75 (2003), No. 4, pp. 259–274.